

The Arecibo OH Megamaser Survey and the Galaxy Merger Rate

Jeremy Darling & Riccardo Giovanelli

Department of Astronomy, Cornell University, Ithaca NY, 14853

Abstract. We present the current results of a survey for OH megamasers (OHMs) underway at the Arecibo Observatory¹. The survey is 2/3 complete and has produced a high OHM detection rate (1 in 6) from a redshift-selected sample of *IRAS* galaxies. The survey will relate the OHM luminosity function to the galaxy merger rate, allowing subsequent blind OHM surveys to measure the galaxy merger rate as a function of cosmic time. The survey has also made the first detection of strong variability in OHMs. Variability will provide a powerful tool for understanding the small-scale physical settings and mechanisms of OHMs.

1. The Survey

The Arecibo OH megamaser (OHM) survey selects candidates from the PSCz redshift catalog (Saunders et al. 2000) with the criteria: (1) $f_{60\mu m} > 0.6$ Jy, (2) $0.1 < z < 0.45$, and (3) $0^\circ < \delta < 37^\circ$ (Darling & Giovanelli 2000). With a detection rate of 1 OHM in 6 candidates, the complete survey will double the sample of OHMs to roughly 100 objects. The survey has identified 35 new OHMs in luminous infrared galaxies to add to the sample of 55 found in the literature. There is a strong bias for the most FIR-luminous galaxies to host OHMs, and a weak FIR color dependence (see Figure 1). OHMs are detectable out to $z = 3-5$ with modern instruments, and can thus be used to measure the high luminosity tail of the luminous IR galaxy luminosity function for redshifts spanning the epoch of major galaxy mergers ($0.5 < z < 5$). Blank field surveys for OHMs at various redshifts can also measure the galaxy merger rate as a function of cosmic time (Briggs 1998).

Variability has been detected in several OHMs, and is currently under investigation. The variability appears over time scales of months in individual spectral features rather than in broad-band modulation which could be attributed to antenna calibration or pointing errors. Variability in OHMs constrains the sizes of the variable and quiescent spectral features, regardless of the source of modulation (intrinsic to the source or due to propagation effects). Intrinsically variable regions would have sufficiently small angular sizes that they would also be expected to scintillate (see Walker 1998). We thus attribute the variabil-

¹The Arecibo Observatory is part of the National Astronomy and Ionosphere Center, which is operated by Cornell University under a cooperative agreement with the National Science Foundation.




Figure 1. Observed OH Megamaser Candidates. The left panel shows L_{FIR} versus FIR color for 197 candidates observed to date, and the right panel shows the OHM fraction as a function of L_{FIR} . Points with error bars are non-detections at $100\mu m$. Vertical error bars indicate the possible range of L_{FIR} , constrained by $f_{60\mu m}$ and an upper limit on $f_{100\mu m}$. Horizontal arrows indicate upper limits on FIR color. Inset percentages indicate the OHM fraction for each sector delineated by the dashed lines. The inset numbers follow the key: N = Observed (OHMs, Non-Detections).

ity to interstellar scintillation, which gives a weaker constraint on the sizes of emission regions than intrinsic variability. Variability in OHMs, particularly those with $0.1 < z < 1.0$, will provide a powerful tool for understanding the small-scale physical settings and mechanisms of masers which can be observed at cosmological distances.

Acknowledgments. The authors are very grateful to Will Saunders for access to the PSCz catalog and to the excellent staff of NAIC for observing assistance and support. This research was supported by Space Science Institute archival grant 8373 and made use of the NASA/IPAC Extragalactic Database (NED) which is operated by the Jet Propulsion Laboratory, California Institute of Technology, under contract with the National Aeronautics and Space Administration.

References

- Briggs, F. 1998, *A&A*, 336, 815
 Darling, J. & Giovanelli, R. 2000, *AJ*, 119, 3003
 Saunders, W., et al. 2000, in *Cosmic Flows: Towards an Understanding of the Large-Scale Structure in the Universe*, ed. S. Courteau, M. Strauss & J. Willick (San Francisco: ASP), in press
 Walker, M. A. 1998, *MNRAS*, 294, 307